

2025-26 COVID-19 Vaccine Routine Vaccination Schedule

Age	Vaccine	If unvaccinated:	If had any prior doses, give 2025-26 doses:
6 – 23 months	Moderna–Spikevax (pediatric)		<p>If 1 prior Moderna dose: 4-8 weeks → 1</p> <p>If 1 prior Pfizer dose: 4-8 weeks → 1 ≥8 weeks → 2</p> <p>If ≥2 prior Moderna/Pfizer doses: ≥8 weeks → 1</p>
2 – 4 years ^{^^}	Moderna–Spikevax (pediatric)	1 Dose	<p>If ≥1 prior Moderna/Pfizer dose: ≥8 weeks → 1</p>
5 – 11 years ^{^^}	Moderna–Spikevax (pediatric)	1 Dose	<p>If 1 or more prior doses (of any of the brands), then[^]: ≥2 months → 1 Spikevax/Comirnaty</p>
	Pfizer–Comirnaty (pediatric)	1 Dose	
12+ years ^{^^}	Pfizer–Comirnaty (adol/adult)	1 Dose	<p>If 1 or more prior doses (of any of the brands), then[^]:</p> <p>Ages 12–64 years^{^^}: ≥2 months[§] → 1 Spikevax/Mnexspike/Comirnaty/Nuvaxoid</p> <p>Ages 65+ years: ≥2 months[§] → 1 6 months[§] → 2</p>
	Moderna–Spikevax (adol/adult)	1 Dose	
	Moderna–Mnexspike	1 Dose	
	Sanofi/Novavax Nuvaxoid	1 Dose	

[^] Janssen (J & J) vaccine has been deauthorized. Follow schedule for 12+ years for any prior doses.

[§] Minimum interval 2 months, except for Mnexspike, the minimum interval is 3 months.

^{^^} Vaccination is especially recommended to 2-64 year olds who have risk factors, are in close contact with others with risk factors, or have never been vaccinated.

Adapted from the California Department of Public Health

2025-26 COVID-19 Vaccine Moderately/Severely Immunocompromised Vaccination Schedule

Age	Vaccine	If unvaccinated:	If had any prior doses, give 2025-26 doses:
6 months–4 years	Moderna–Spikevax (pediatric)	1st Dose → 4 weeks → 2nd Dose → ≥4 weeks → 3rd Dose → 6 months [§] → Additional Dose(s)*	If prior doses of Moderna or Pfizer at 6m-4y: 1 prior dose: 4 w → 1 → ≥4 w → 2 → 6 m [§] → Additional Dose(s)* 2 prior doses: → ≥4 w → 1 → 6 m [§] → Additional Dose(s)* ≥3 prior doses**: → ≥8 w → 1 (for ages 5+ yrs, Pfizer dose is also OK)
	Moderna–Spikevax (pediatric)	1st Dose → 4 weeks → 2nd Dose → ≥4 weeks → 3rd Dose → 6 months [§] → Additional Dose(s)* Spikevax/Comirnaty	1 prior dose: 3 w → 1 → ≥4 w → 2 → 6 m [§] → Additional Dose(s)* 2 prior doses: → ≥4 w → 1 → 6 m [§] → Additional Dose(s)* ≥3 prior doses**: → ≥8 w → 1
5–11 years	Pfizer–Comirnaty (pediatric)	1st Dose → 3 weeks → 2nd Dose → ≥4 weeks → 3rd Dose → 6 months [§] → Additional Dose(s)* Spikevax/Comirnaty	1 prior dose: 3 w → 1 → ≥4 w → 2 → 6 m [§] → Additional Dose(s)* 2 prior doses: → ≥4 w → 1 → 6 m [§] → Additional Dose(s)* ≥3 prior doses**: → ≥8 w → 1
	Pfizer–Comirnaty (adol/adult)	1st Dose → 3 weeks → 2nd Dose → ≥4 weeks → 3rd Dose → 6 months [§] → Additional Dose(s)* Spikevax/Comirnaty/Nuvaxoid/Mnexspike	1 prior dose: 4 w → 1 → ≥4 w → 2 → 6 m [§] → Additional Dose(s)* 2 prior doses: → ≥4 w → 1 → 6 m [§] → Additional Dose(s)* ≥3 prior doses**: → ≥8 w → 1
	Moderna–Spikevax (adol/adult)	1st Dose → 4 weeks → 2nd Dose → ≥4 weeks → 3rd Dose → 6 months [§] → Additional Dose(s)* Spikevax/Comirnaty/Nuvaxoid/Mnexspike	1 prior dose: 4 w → 1 → ≥4 w → 2 → 6 m [§] → Additional Dose(s)* 2 prior doses: → ≥4 w → 1 → 6 m [§] → Additional Dose(s)* ≥3 prior doses**: → ≥8 w → 1
	Moderna–Mnexspike	1st Dose → 4 weeks → 2nd Dose → ≥4 weeks → 3rd Dose → 6 months [§] → Additional Dose(s)* Spikevax/Comirnaty/Nuvaxoid/Mnexspike	1 prior dose: 4 w → 1 → ≥4 w → 2 → 6 m [§] → Additional Dose(s)* 2 prior doses: → ≥4 w → 1 → 6 m [§] → Additional Dose(s)* ≥3 prior doses**: → ≥8 w → 1
12+ years	Sanofi/Novavax Nuvaxovid	1st Dose → 3 weeks → 2nd Dose [¶] → 6 months [§] → Additional Dose(s)* Spikevax/Comirnaty/Nuvaxoid/Mnexspike	1 prior dose: → ≥3 w → 1 → 6 m [§] → Additional Dose(s)* ≥2 prior doses**: → ≥8 w → 1
	Sanofi/Novavax Nuvaxovid	1st Dose → 3 weeks → 2nd Dose [¶] → 6 months [§] → Additional Dose(s)* Spikevax/Comirnaty/Nuvaxoid/Mnexspike	1 prior dose: → ≥3 w → 1 → 6 m [§] → Additional Dose(s)* ≥2 prior doses**: → ≥8 w → 1

* Further doses may be given under shared clinical decision-making at a minimum interval of 2 months.

** Ages 5-11 years may be given Moderna or Pfizer after ≥3 prior doses. Ages 12+ years may be given Spikevax, Comirnaty, Nuvaxovid, or Mnexspike.

§ Minimum interval 2 months, except for Mnexspike, the minimum interval is 3 months.

¶ If >8 weeks passed since the first Novavax dose, any 2025–26 COVID-19 vaccine (Spikevax/Comirnaty/Nuvaxovid/Mnexspike) may be given.

Adapted from the California Department of Public Health