


Counties: Appanoose, Black Hawk, Cass, Cerro Gordo, Clinton, Des Moines, Fremont, Harrison, Henry, Jefferson, Lee, Mahaska, Marshall, Monona, Monroe, Montgomery, Muscatine, Page, Pottawattamie, Scott, Wapello, Webster and Woodbury

Evidence-based Home Visiting Models: Healthy Families America (HFA), Nurse Family Partnership (NFP), Parents as Teachers (PAT)

68,857 Children ages 0-5

10,561 Births in 2024



U.S. Census Bureau, ACS 5 Year Estimates, Table B09001 - Population Under 18 Years by Age
Iowa Vital Statistics Report 2024, Table 48

Fiscal Year 2025 (FY25) Data


852 Households served

530 Continuing families

322 Newly enrolled

142 Prenatally enrolled

12,414 Total Visits



Priority Characteristics

Low-income Household: 52%

Percentage of participants in households under 100% of the Federal Poverty Level (FPL).

Teenage Pregnancy: 4%

Percentage of parents enrolled in home visiting with a pregnancy who are under 21 years old.

Education: 18%

Low student achievement in the household.

Military: 4%

Current or former military service in the household.

Substance Use: 22%

Parents with a history of substance abuse.

Tobacco: 19%

Tobacco use in the home.

Abuse: 21%

History of child maltreatment or involvement with Child Protective Services (CPS).


Disabilities: 19%

Child with developmental delays or disabilities in the household.

Outcomes


Breastfeeding at 6 Months

Infants born to mothers who enrolled prenatally and received breastmilk of any amount at 6 months of age.




Depression Screening

Caregivers screened for depression using the EPDS within 3 months of enrollment or delivery.




Postpartum Care

Mothers enrolled prenatally or within 30 days after delivery who received a postpartum visit within 8 weeks of delivery.




Tobacco Cessation Referral

Primary caregivers who reported using tobacco at enrollment, not in tobacco cessation services, who were referred to tobacco cessation services within 3 months of enrollment.




Safe Sleep

Infants enrolled in home visiting that were always placed to sleep on their back without bedsharing or soft bedding.



Parent-Child Interaction Screening


Primary caregivers who received a caregiver-child interaction observation by the home visitor using the PICCOLO, CHEERS, or DANCE.



Outcomes


Early Language and Literacy Activities

Target children whose caregiver reported that during a typical week, they read, told stories and/or sang songs with their child every day.




Developmental Screening

Children who were screened on-time for development delays using the ASQ-3 at 9 months, 18 months and 30 months.




Intimate Partner Violence Screening

Primary caregivers screened within 6 months of enrollment using the Relationship Assessment Tool.



Continuity of Insurance Coverage

Primary caregivers who had continuous health insurance coverage for at least 6 consecutive months in the year.



Completed Depression Referrals

Primary caregivers referred to services for a positive screen for depression who received at least one service contact from a provider.

