

Immunization Schedule with Combination Vaccines

	2 MONTHS	4 MONTHS	6 MONTHS	12 MONTHS	15 MONTHS	18 MONTHS	4-6 YEARS		
PEDIARIX® PROQUAD® QUADRACEL™ or KINRIX®	PEDIARIX® DTaP, IPV, HepB + PCV Rotavirus Hib	PEDIARIX® DTaP, IPV, HepB ¹ + PCV Rotavirus Hib	PEDIARIX® DTaP, IPV, HepB + PCV Rotavirus ² Hib ³	HepA MMR ⁴ Varicella ⁴ PCV ⁵ Hib ⁵	DTaP	HepA	QUADRACEL™ or KINRIX®⁷ DTaP, IPV + PROQUAD® MMRV		
	PENTACEL®⁵ PROQUAD® QUADRACEL™ or KINRIX®	PENTACEL® DTaP, IPV, Hib + PCV Rotavirus HepB	PENTACEL® DTaP, IPV, Hib + PCV Rotavirus HepB ¹	PENTACEL® DTaP, IPV, Hib + PCV Rotavirus ² HepB	HepA MMR ⁴ Varicella ⁴ PCV ⁵	PENTACEL® DTaP, IPV, Hib	HepA	QUADRACEL™ or KINRIX®⁷ DTaP, IPV + PROQUAD® MMRV	
		VAXELIS™⁶ PROQUAD® QUADRACEL™ or KINRIX®	VAXELIS™ DTaP, IPV, Hib, HepB + PCV Rotavirus	VAXELIS™ DTaP, IPV, Hib, HepB ¹ + PCV Rotavirus	VAXELIS™ DTaP, IPV, Hib ³ , HepB + PCV Rotavirus ²	HepA MMR ⁴ Varicella ⁴ PCV ⁵ Hib ⁵	DTaP	HepA	QUADRACEL™ or KINRIX®⁷ DTaP, IPV + PROQUAD® MMRV

Everyone 6 months+: 1) COVID-19 vaccine per current recommendations 2) Flu vaccine every fall

1st RSV SEASON: RSV immunization for all infants < 8 months.⁸

*Adapted from the California Department of Health Immunization Branch

Make sure the vaccine you administer Keep it simple. Stick with the same product.

This is a suggested schedule for VFC providers ordering combination vaccines. For alternatives and details, consult the latest recommended [Immunization Schedules](#) from CDC.

1. A dose of Hepatitis B vaccine is not necessary at 4 months if doses are given at birth and 2 months but may be included as part of a combination vaccine.
2. The six month dose is not needed if Rotarix[®] was used exclusively for both dose 1 and 2 of the rotavirus vaccine series.
3. This six month Hib dose is not indicated if PedvaxHIB[®] is used exclusively for the 2 and 4 month infant doses.
4. CDC recommends MMR + Varicella at 12-15 months. Providers can use their discretion whether to use MMRV, however.
5. Can be administered as late as 15 months. For more information, consult the [Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger](#), United States, 2023.
6. Licensed by FDA for children 6 weeks through 4 years of age (prior to the 5th birthday).
7. Licensed by FDA for children 4 through 6 years with previous doses of INFANRIX[™] or PEDIARIX[™]. ACIP recommends that, whenever feasible, the same manufacturer's DTaP vaccines be used for each dose in the series; however, vaccination should not be deferred because the type of DTaP previously administered is unavailable or unknown. See www.cdc.gov/mmwr/preview/mmwrhtml/mm5739a4.htm.
8. Infants <8 months entering their first RSV season should receive RSV immunization if birth parent's prenatal RSV vaccination status is: unvaccinated, unknown, or vaccinated <14 days before birth. RSV immunization is also recommended for specified children ages 8-19 months who are at increased risk of severe RSV and entering their 2nd RSV season.

Iowa HHS Vaccines for Children Program 1-800-831-6293