



## Vaccines for Children (VFC) Program Available Vaccines and Covered Age Ranges

The following vaccines are available through the Iowa Vaccines for Children (VFC) Program. VFC-eligible children birth through 18 years of age must meet one of the following eligibility criteria: Medicaid, uninsured, American Indian/Alaskan Native, or underinsured. Underinsured children (health insurance but the benefit plan does not include immunizations) are eligible for VFC vaccine at Federally Qualified Health Centers (FQHC), Rural Health Centers (RHC), and local public health agencies.

Diseases and ACIP-Recommended Vaccines Covered by the Iowa VFC Program		
Vaccine Group/Disease	Vaccine(s) and other immunizing agents	Covered Ages for VFC-eligible Children
Chickenpox (Varicella)	Varivax®	12 months through 18 years of age
COVID-19	Comirnaty®	12 through 18 years of age
	Pfizer-BioNTech COVID-19 Vaccine (6m-4Y)	6 months through 4 years of age
	Pfizer-BioNTech COVID-19 Vaccine (5-11Y)	5 through 11 years of age
	Spikevax™	12 through 18 years of age
	Moderna COVID-19 Vaccine (6m-11Y)	6 months through 11 years of age
	Novavax COVID-19 Vaccine	12 years through 18 years of age
DTaP and DTaP Combination	DTaP (Daptacel® & Infanrix®) <sup>1</sup>	6 weeks through 6 years of age
	DTaP-IPV (Kinrix & Quadracel) <sup>5</sup>	4 through 6 years of age for the fifth dose in the DTaP series and the fourth dose in the IPV series only
	DTaP-IPV-Hib (Pentacel®) <sup>6</sup>	6 weeks through 4 years of age
	DTaP-IPV-HepB (Pediatrix®) <sup>7</sup>	6 weeks through 6 years of age for the first three doses of the primary series of DTaP
	DTaP-IPV-Hib-HepB (Vaxelis™) <sup>8</sup>	6 weeks through 4 years of age for the first 3 doses of the primary series of DTaP, IPV, Hib. It should not be used for the HepB birth dose (4 doses of HepB are permitted when combination vaccine is used after the birth dose)
Hib (Haemophilus influenzae type b)	PedvaxHib®, ActHib®, Hiberix®	6 weeks through 59 months of age. Also available for VFC eligible children 5 through 18 years of age with certain underlying medical conditions
Hepatitis A	Vaqta® & Havrix®	1 through 18 years of age
Hepatitis B	Engerix B® & Recombivax HB®	Birth through 18 years of age. Birthing Hospitals only: Birth dose vaccine available for entire birth cohort
Human Papillomavirus (HPV)	Gardasil®9	9 through 18 years of age
Influenza (Flu)		6 months through 18 years of age

Meningococcal and Meningococcal Combination	MenACWY (MenQuadfi™ & Menveo®)	11 through 18 years of age. Also available for VFC eligible children aged 2 months through 10 years of age who are at increased risk for developing meningococcal disease
	MenB (Trumenba® & Bexsero®)	16 through 18 years of age. Also available for VFC children 10 through 18 years of age who are at increased risk due to a medical condition or a Serogroup B disease outbreak.
	MenABCWY (Penbraya™)	16-18 years when both MenACWY and MenB are indicated at the same visit. Also available for children 10-18 years who are at increased risk for meningococcal disease attributable to serogroups A,C,W,Y and B.
Measles, Mumps, Rubella (MMR) and MMR Combination	MMR (MMR®II & Priorix) <sup>3</sup>	12 months through 18 years of age
	MMRV (Proquad®) <sup>4</sup>	12 months through 12 years of age
Pneumococcal	PCV15 (Vaxneuvance™)	6 weeks through 59 months of age. Also available for children 5 through 18 years with certain underlying medical conditions who are at increased risk for invasive pneumococcal disease
	PCV20 (Prevnar 20™)	6 weeks through 59 months of age. Also available for children 5 through 18 years with certain underlying medical conditions who are at increased risk for invasive pneumococcal disease
	PPSV23 (Pneumovax®23)	2 through 18 years of age. Must obtain prior approval from the Iowa VFC Program
Polio (IPV)	IPOL®	6 weeks through 18 years of age
Rotavirus	Rotarix® & RotaTeq®	6 weeks to 8 months 0 days of age
Respiratory Syncytial Virus (RSV)	Monoclonal Antibody (Beyfortus™) – limited availability	<8 months born during or entering their 1 <sup>st</sup> RSV season. Also available for children 8-19 months at increased risk for severe RSV disease entering their 2 <sup>nd</sup> RSV season
	Maternal RSV vaccine (Abrysvo™) – limited availability	Pregnant individuals 18 years of age or younger at 32 through 36 weeks gestational age
Tetanus (Td and Tdap Combination vaccines)	Td preservative-free (TDVAX & Tenivac®) <sup>1</sup> – limited availability	4 months through 18 years of age. Must obtain prior approval from the Iowa VFC Program
	Tdap (Adacel® & Boostrix®) <sup>2</sup>	7 through 18 years of age
<sup>1</sup> DTaP and Tdap combine protection against diphtheria, tetanus, and pertussis <sup>2</sup> Td combines protection against diphtheria and tetanus <sup>3</sup> MMR combines protection against measles, mumps, and rubella <sup>4</sup> MMRV is a combination vaccine containing MMR and varicella <sup>5</sup> Kinrix and Quadracel are combination vaccines containing DTaP and IPV <sup>6</sup> Pentacel is a combination vaccine containing DTaP, IPV, and Hib <sup>7</sup> Pediarix is a combination vaccine containing DTaP, IPV, and HepB <sup>8</sup> Vaxelis is a combination vaccine containing DTaP, IPV, Hib, and HepB  Source: Centers for Disease Control and Prevention (CDC)		